

University of Groningen

Is begging cheap?

Verhulst, S; Wiersma, P

Published in:
The Auk

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1997

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Verhulst, S., & Wiersma, P. (1997). Is begging cheap? *The Auk*, 114(1), 134-134.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Is Begging Cheap?

SIMON VERHULST¹ AND POPKO WIERSMA

Zoological Laboratory, P. O. Box 14, 9750 AA Haren, The Netherlands

Avian nestlings beg to obtain food (von Haartman 1953), and begging is used increasingly as a model to study game-theoretic problems, the evolution of signaling, and parent-offspring conflict (Godfray 1991, Cotton et al. 1996). A fundamental assumption in these studies is that begging is costly (Godfray 1991). McCarty (1996) recently provided the first estimates of the costs of begging. These estimates were obtained by monitoring oxygen consumption of nestlings induced to beg in metabolic chambers. The metabolic rate during begging was increased by 8% in European Starlings (*Sturnus vulgaris*) and by 42% in Tree Swallows (*Tachycineta bicolor*). From these data (in combination with some tentative extrapolations to a nestling's daily energy demands), McCarty concluded that "the energetic cost of begging is surprisingly low" and that "conclusions from models dependent on the assumption of a high cost to begging should be viewed with caution."

However, McCarty expressed costs in energetic terms, whereas the relevant currency in an evolutionary sense reflects the fitness consequences of variations in begging rate. To translate the energetic costs of begging to fitness costs, one needs to know how energy allocation to begging affects growth, as well as the relationship between growth and reproductive prospects of the chick. We believe that such a calculation may lead to a very different conclusion. Among developing birds, only 13–28% of the total metabolized energy is allocated to growth, the remainder being spent largely on maintenance and thermoregulation (Weathers 1992). Thus, a chick spending an extra 10% of its total energy budget on begging may thereby reduce the amount of energy available for growth by 50%. In theory, this problem could be solved by increasing the energy budget (if extra food is obtained with begging), but at some point the energy budget will be constrained by the digestive capacity of the chick (Weiner 1992), or by other factors (Konarzewski et al. 1996). Thus, a small increase in energy expenditure for begging may result in a disproportionate decrease in reproductive prospects of the chick, and it is only this decrease that may yield a relevant estimate of the costs of begging.

Nevertheless, studies of energetics may be critical to bridge the gap between behavior and fitness. In birds, fitness consequences of variation in growth rate are relatively well known (Tinbergen and Boerlijst 1990), and as McCarty (1996) demonstrated, the energetic costs of begging can readily be measured in the laboratory. This makes it feasible to obtain estimates of the link between behavior and fitness, via the energetics of begging and growth, which is rarely possible in free-living animals. Such knowledge in turn will make it possible to develop parameterized models of begging behavior that are sufficiently detailed to be tested against reality.

Acknowledgments.—S.V. is supported by SLW-grant 803-30-165.

LITERATURE CITED

- COTTON, P. A., A. KACELNIK, AND J. WRIGHT. 1996. Chick begging as a signal: Are nestlings honest? *Behavioral Ecology* 7:178–182.
- GODFRAY, H. C. J. 1991. Signalling of need by offspring to their parents. *Nature* 352:328–330.
- KONARZEWSKI, M., J. KOWALCZYK, T. SWIERUBSKA, AND B. LEWONCZUK. 1996. Effect of short-term feed restriction, realimentation and overfeeding on growth of Song Thrush (*Turdus philomelos*) nestlings. *Functional Ecology* 10:97–105.
- MCCARTY, J. P. 1996. The energetic cost of begging in nestling passerines. *Auk* 113:178–188.
- TINBERGEN, J. M., AND M. C. BOERLIJST. 1990. Nestling weight and survival in individual Great Tits (*Parus major*). *Journal of Animal Ecology* 59:1113–1127.
- VON HAARTMAN, L. 1953. Was reizt den Trauerfliegenschnapper zu Futtern? *Vogelwarte* 16:157–164.
- WEATHERS, W. W. 1992. Scaling nestling energy requirements. *Ibis* 134:142–153.
- WEINER, J. 1992. Physiological limits to sustainable energy budgets in birds and mammals: Ecological implications. *Trends in Ecology and Evolution* 7:384–388.

Received 5 August 1996, accepted 21 August 1996.

Associate Editor: T. E. Martin

¹ E-mail: s.verhulst@biol.rug.nl